



High- ℓ Science with PICO

Daniel Green
UC San Diego





High- ℓ Targets

Inflation: $n_s, \alpha_s, \zeta_{\vec{k}}, f_{\text{NL}}^{\text{eq.}}, \dots$

Early universe: Y_p/BBN , isocurvature, $C_{\nu\text{B}}/N_{\text{eff}}$


Dark sectors: forces between SM, DM and/or DR

Other: recombination history, neutron lifetime

High- ℓ Target

N_{eff} is a good reference for high- ℓ science

- Affects damping tail & acoustic structure
 - Has well defined theoretical targets
 - Forecasts do not depend sensitively on low- ℓ
 - Relative improvement similar for other parameters
 - Can be a useful metric for optimizing design
-



$$N_{\text{eff}} \equiv \frac{8}{7} \left(\frac{11}{4} \right)^{4/3} \frac{\rho_\nu}{\rho_\gamma}$$

Perfect decoupling :


$$N_{\text{eff}} = 3.$$


Imperfect decoupling + QED :


$$N_{\text{eff}} = 3.045$$


Salas & Pastor (2016);
Mangano et al. (2005)

Other Light Relics











$$3M_{\text{pl}}^2 H^2 = \rho_\gamma + \rho_\nu$$




$$\frac{1}{k_d^2} \sim \langle \Delta x^2 \rangle \propto t \quad t \propto \frac{1}{H}$$

$$3M_{\text{pl}}^2 H^2 = \rho_\gamma + \rho_\nu$$




$$N_{\text{eff}} = 3.04 \pm 0.18 \text{ Planck (2015)}$$




Phase shift = Free streaming

Bashinsky & Seljak (2003)



$$N_{\text{eff}}^{\text{phase}} = 2.3^{+1.1}_{-0.4}$$

Follin et al. (2015)




$$N_{\text{eff}}^{\text{phase}} = 2.80^{+0.24}_{-0.23}$$

Baumann, DG, Meyers, Wallisch (2015)


PICO Forecasts


Central questions for PICO


- Can PICO reach well-motivated targets?
 - Are there unique advantages to space?
 - Opportunities when combined with ground-based?
 - Does this suggest other interesting observables?
-



w. Meyers, includes delensing from DG, Meyers, van Engelen







DG forecasts, do not include delensing

Take-away


PICO would have comparable sensitivity to CMB-S4:

- CMB-S4 beam reaches higher ℓ
 - PICO has larger f_{sky}
 - These two roughly offset
 - Lower PICO noise limited because of CV
-


Y_p marginalization

PICO measures more E-modes with $\ell \lesssim 2000$

- Acoustic peaks are sharper in E-modes
- Can lead to better constraints
- The neutrino phase shift is one such observable
- Capture this effect by marginalizing over Y_p



Baumann, DG, Meyers, Wallisch (2015)




DG forecasts, do not include delensing

Lensing & Delensing (T/E)

Improvements from CMB (de)lensing will help more:

- EB estimator can benefit from lower noise of PICO
- Delensing can improve parameter constraints



DG, Meyers & van Engelen (2016)

Summary

High- ℓ science with PICO driven by f_{sky}

- More CVL modes, especially E-modes
 - Ground benefits most from high- ℓ TE
 - Combining PICO and CMB-S4 is useful
-



Thank you