

# Welcome to the PICO Workshop

---

- WIFI
  - use the 'eduroam' network
  - if you don't have your account, look at the back of your name tag
  - Use 'username@umn.edu' + provided password.
- Dinners (both walking distance, please sign dinner sheet)
  - today 6:30 pm Cafe421 (Italian/American, white tablecloths)
  - tomorrow 6:30 pm Bona (Vietnamese, no tablecloths)
- Poster session tomorrow
- Please upload your talks
- Please stay on time
- Discussions: leave enough time for audience participation
- If you are not on the microphone - webexers can not hear you



# PICO- Context, Design, Capabilities, Workshop

Shaul Hanany  
PICO Workshop, May 2018



## 2010 Decadal Panel: New space Activities - Medium Projects

---

- CMB listed as a strategic program (priority 2, after exoplanet searches)
- Sub-orbital program to continue search for the B-mode signal
- Continued investment in technology development
- “A successful detection of B-modes from inflation could trigger a mid-decade shift in focus toward preparing to map them over the entire sky.”
- To paraphrase: wait and see what we learn from Planck



# 2010 Astrophysics Decadal Panel Recommendations

---

- Flagship: WFIRST (>\$1B)
- Explorer
  - Missions of Opportunity \$65M
  - Small Explorer ~\$150M
  - Explorer \$250M
- No Probe-class astrophysics missions with \$250 - \$1000M
- Planetary Science Decadal Survey
  - New Frontier-class missions: <=\$1000M
  - Juno (2011); OSIRIS-REx (2016); now competing next

EPIC - IM



# NASA's Preparations for 2020

---

- **Set up 8 Probe Mission Studies; Probe= \$400M-\$1000M**
  - Transient Astrophysics Probe (Camp, GSFC)
  - Cosmic Dawn Intensity Mapper (Cooray, UC Irvine)
  - Cosmic Evolution through UV Spectroscopy (Danchi, GSFC)
  - Galaxy Evolution Probe (Glenn, UColorado)
  - **Inflation Probe (Hanany, University of Minnesota)**
  - High Spatial Resolution X-Ray Probe (Mushotzky, UMaryland)
  - Multi-Messenger Astrophysics (Olinto, UChicago)
  - Precise Radial Velocity Observatory (Plavchan, Missouri State)



# Study Outcomes

---

- Studies will produce 50 pg. reports + cost estimates that will be submitted to NASA and to the Decadal Panel (12/2018)
- Possible outcomes:
  - Panel recommends a funding wedge. Probes are competed later.
  - Panel recommends specific missions
  - Any combination of the above
- Our desired outcome: panel recognizes the promise of a future space mission and gives high ranking



# PICO Study

---



- Phase 1: decide on science goals and overall design drivers
  - Spectrometer, Imager, or both; resolution, survey type, frequency coverage, depth
- Phase 2:
  - instrument + mission design and costing;
  - assess performance
- Phase 3: write report
- Submit report by end of December 2018





# Spectrometer, Imager, or both?

- Which instrument(s) to implement?
  - Considered superPIXIE = x10 PIXIE sensitivity
  - Considered combined PIXIE + Imager
- EC conclusion: there is strong case for two separate missions, one devoted to spectroscopy and another to imaging (or a more expensive single mission)
- A combined mission within \$1B cap will weaken both instruments
- Design and costing exercise will concentrate on an imager.
- Paradigm: design the most scientifically compelling mission within the cost cap





# Design Basics



---

- Full sky
  - Inflation including the reionization peak
  - Star formation history
- Broad frequency coverage
  - Galactic emissions (on their own and for foregrounds)
- Resolution
  - Neff
  - Inflation, neutrino mass (through lensing potential and delensing)
  - Galactic science




# PICO in Brief

- Millimeter/submillimeter-wave, polarimetric survey of the entire sky
- 21 bands between 20 GHz and 800 GHz
- 1.4 m aperture telescope
- Diffraction limited resolution: 38' to 1'
- 13,000 transition edge sensor bolometers + multiplexed readouts
- 5 year survey from L2
- $0.6 \text{ uK} \cdot \text{arcmin}$  (*Planck* = 50 ; S4 =  $0.8 \text{ uK} \cdot \text{arcmin}$ , 3%)










# PICO Science

- Inflation:  $r < 10^{-4}$  (95%),  $\sigma(r) = 5 \cdot 10^{-5}$
- Cosmic variance limited  $\tau$ ,  $\sigma(\tau) = 0.002$
- Neutrino mass:  $\sigma(\Sigma m_\nu) = 14$  meV (inc. DASI BAO; equivalent independent limit from cluster counts)
- $\sigma(N_{eff}) = 0.03$
- $\sigma(\omega_0) = 0.023$  ,  $\sigma(\omega_a) = 0.13$  with 140,000 clusters
- Correlate lensing map with other mass tracer surveys










# PICO Science Goals

- Map magnetic field over 70% of galaxy with 1' resolution
- Resolve B-field structure in 8 nearby clouds on core scale (0.1 pc) (currently none)
- Resolve B-field structure in 2000 clouds with 1 pc resolution to compare roles of turbulence and B-field in star formation efficiency (currently 14)
- Map sub-mm emission in ISM of 70 nearby galaxies (handful to date)

■ PICO, 1'   
 ■ Planck, 5'   
 ■ PICO, 5'








# PICO Science Goals

- Discover 3000 highly magnified dusty galaxies at  $z$  up to  $\sim 4.5$ ;
- Discover 3000 proto-clusters over the sky and extending to high redshift;
- Detect polarization of 4000 radio and FIR-emitting galaxies;
- $\times 10$ -100 more than known today
- Probe star formation history; determine galaxy and cluster formation and evolution; learn about dark matter substructure; and measure properties of jets in radio-loud sources.



# Instrument Design



- 2 mirror, Open-Dragone, 1.4 aperture telescope
- Primary at ambient temperature
- 4-6 K-cooled Stop + Secondary
- 0.1 K focal plane





# Focal Plane

- 3 color antenna coupled for pixels A - F  $\nu \leq 500$  GHz
- Single color, horn-coupled, absorber-based pixels for G,H,I  $\nu > 500$  GHz
- All based on TES bolometers
- Total bolometer count = 12,996
- Multiplexed readouts (TDM: x128 columns x 102 rows)
- Alternating columns are oriented as Q / U pixels




|   |              |
|---|--------------|
| A | 21, 30, 43   |
| B | 25, 36, 52   |
| C | 62, 90, 129  |
| D | 70, 108, 155 |

|         |               |
|---------|---------------|
| E       | 186, 268, 385 |
| F       | 223, 321, 462 |
| G, H, I | 555, 666, 799 |



# Orbit + Scan



- Orbit around L2
- $\alpha \sim 25$ ,  $\beta \sim 70$
- Spin = 1 rpm
- Precess = 10 hours
- ~6 months for single full sky survey
- 5 year total survey




Each circle is one spin •

OD =  $\alpha + \beta$  •

ID =  $\alpha - \beta$  •







# PICO Summary

- Inflation, quantum gravity, particle physics, extragalactic and galactic structure and evolution:
  - All unique goals for the PICO measurements
  - PICO is the only instrument with the combination of sky coverage, resolution, frequency bands, and sensitivity to achieve all of this science with one platform.
- Initial engineering + costing study complete:
  - Technology implementation is a simple extension of today's technologies; no technological breakthroughs required
  - Mission is a good fit to the cost window



# PICO Study

---

- Phase 1: decide on science goals and overall design drivers
  - Spectrometer, Imager, or both; resolution, survey type, frequency coverage, depth
- Phase 2:
  - instrument + mission design and costing: Completed two TeamX studies + costing
  - assess performance
- Phase 3: write report
- Submit report by end of December 2018





# Workshop Goals

---

- Review where we are, decide what else to do for the report, discuss what goes into the report, distribute writing assignments
- File with candidate 'Issues to Discuss' posted
- Day 1, Science:
  - What classes of models is it compelling to rule out in addition to and at levels of  $r$  below the Starobinsky-type class of models?
  - Which science goals have we not included yet? What other science targets that are well suited for space have we not yet thought about?
  - Ancillary science: are we properly reaching out to other communities?
  - Is there a science argument for a Guest Observer program?
  - How are the various science goals affected by foregrounds?
- Throughout: how do PICO and S3/S4 complement and strengthen each other?



Additional Slides




# LiteBIRD

---



- It would have been reasonable to consider a smaller aperture mission
- It would have been reasonable to consider how such smaller aperture mission complements S4
- A smaller aperture mission (a-la LiteBIRD) is more limited in its science capabilities (Galactic science, Cluster science, Lensing (=neutrino mass) and Neff)
- A US-based smaller aperture mission is not much cheaper than PICO
- LiteBIRD is in a proposal stage
- We saw our task as putting forth the most scientifically compelling mission that is safely within the cost window






# Simple Foreground Model

- 2 component dust model (a-la Finkbeiner et al)
- Synchrotron with power law frequency dependence
- $\ell$  dependence consistent with Planck and WMAP
- Includes correlation between dust and synchrotron, consistent with current data
- Model does not include:
  - spatial variation of the spectral index
  - spatial variation of dust temperature
- Foreground separation based on ILC
- 40% of sky (70% of sky reduces  $\sigma(r)$ )







# Delensing

- Iterative delensing post-ILC foreground separation
- Lensing reduction by a factor of  $\sim 7$ :  $A_L = 0.14$
- $S/N > 10$  on lensing potential power spectrum across broad range of  $\ell$





Figure: R. Flauger

Figure: R. Flauger





Figure: R. Flauger





# Balloons

---

- Currently funded: PIPER, SPIDER (2nd flight)
- Proposed: IDS, BFORE, Dust Buster
- IDS:
  - 20,000 detectors,
  - 7 bands, 150 - 360 GHz
  - Combined observations with BICEP/Keck (10 bands total)




# What's at Stake

---

- NASA only invests in technology development or balloon payloads that lead to a future space mission.
- Over the years NASA has spent significant resources in CMB activities (space, balloons, tech development) because there was a mission in the future.
- NASA invests only in what the decadal panel recommends
- Many of us (most? all?) recognize the strengths of a future CMB space mission, the complementarity with sub-orbital, and the value of keeping NASA engaged with CMB






# S4 Inflation Constraints

- Designed to provide detection of  $r > 0.003$
- $r < 0.001$  (95%)
- 3-8% of sky
- $r \geq 0.004$  (5 sigma) in 4 years
- $r \geq 0.003$  (5 sigma) in 8 years



# Optics + Cooling

- Open Dragone Telescope
  - No direct view to sky
  - No three-reflection sidelobe
  - Cold stop (without cooling primary mirror)
- Design includes enhancement to DLFOV through coma correction
- Primary mirror at  $\sim 40$  K;
- Stop + secondary actively cooled to  $\sim 6$  K;
- Focal plane @  $0.1$  K with cADR






# *LiteBIRD*


Masashi Hazumi

- 1) Institute of Particle and Nuclear Studies (IPNS), High Energy Accelerator Research Organization (KEK)
- 2) Kavli Institute for Mathematics and Physics of the Universe (Kavli IPMU), The University of Tokyo
- 3) Graduate School for Advanced Studies (SOKENDAI)
- 4) Institute of Space and Astronautical Science (ISAS), Japan Aerospace Exploration Agency (JAXA)






“Current Status  
of LiteBIRD in  
JAXA” by Toru  
Yamada  
(Former ISAS  
Director of  
International  
Strategy and  
Coordination)







## Full success of LiteBIRD

- $\sigma(r) < 1 \times 10^{-3}$  (for  $r=0$ )
- All sky survey (for  $2 \leq \ell \leq 200$ )\*

### Remarks

1.  $\sigma(r)$  is the total uncertainty on the  $r$  measurement that includes the following uncertainties\*\*
  - statistical uncertainties
  - instrumental systematic uncertainties
  - uncertainties due to residual foregrounds and bias
  - uncertainties due to lensing B-mode
  - cosmic variance (for  $r > 0$ )
  - observer bias
2. The above should be achieved without delensing.





\* **More precise (i.e. long) definition ensures  $>5\sigma$   $r$  detection from each bump for  $r > 0.01$ .**

\*\* We also use an expression  $\delta r = \sigma(r=0)$ , which has no cosmic variance.




# LiteBIRD

- $\sigma(r) < 0.001$  (for  $r=0$ )
- Imager, 35 - 450 GHz
- 15 frequency bands (some overlap)
- Two telescopes; 0.5 deg resolution at 100 GHz
- $2.5 \mu\text{K}\cdot\text{arcmin}$  in polarization






# LiteBIRD U.S. Deliverables (A. LEE, US PI)




HF = High Frequency, MF = Mid Frequency, LF = Low Frequency




to Europe

Sub-Kelvin Cooler




230 mm

HF-FPU



LF/MF-FPU




SQUID Amplifiers




150 mm


HF Module




LF Module



MF Module








- Japan:
  - Phase A1 will conclude in 8/2018
  - Then downselect
- US:
  - 2016 Mission of Opportunity (\$65M) proceeded to PhaseA; PhaseB declined
  - Technology development continues
  - Will submit at next MO (2019?)








# PICO and Sub-Orbital CMB Efforts

PICO's capabilities are not matched by any other foreseeable experiment

- Full sky coverage with  $\sim 4'$  resolution (and the same depth S4 has on 5% of the sky)
- Access to the entire range of angular scales of the B-mode signal, including the largest, while maintaining the capability to delens







# PICO and Sub-Orbital CMB Efforts

- Unmatched/unmatcheable frequency coverage
- Galactic foregrounds are known to overwhelm the cosmological B-mode signal
- Signals are at the nano-K level: even low level of residual foregrounds can bias the measurement
- Space gives the most systematic-error-robust platform
- Signals are at the nano-K level

