

Love Wave Formalism

Levi Walls

June 15, 2016

0.1 SH-waves

Another class of surface waves worth considering are horizontally polarized shear (SH) waves also known as Love waves. Considering a single SH-wave propagating in the $\hat{\Omega}$ direction with particle displacement perpendicular and horizontally polarized with respect to the line of propagation, the analysis follows as above. We begin with the displacement field modified from Lay and Wallace (1995), Aki and Richards (2009):

$$\vec{l}(\vec{x}, t) = l_1 \cos(\vec{k} \cdot \vec{x} - \omega t) \vec{e}_H(\hat{\Omega}), \quad (1)$$

where the naming conventions follow that of shear waves presented earlier; here $A = H$ to signify the horizontal polarization of the wave. For the sake of argument, we assume a frequency-independent eigenfunction solution: $l_1 = e^{-z/h}$. This is done to illustrate the attenuation of these waves as one goes further underground. The nature of these eigenfunction solutions is discussed briefly in subsection 0.1.1. Furthermore, the plane-wave expansion of this displacement field is:

$$\vec{l}(\vec{x}, t) = \int df d\hat{\Omega} e^{-z/h} S_H(f, \hat{\Omega}) \vec{e}_H(\hat{\Omega}) e^{2\pi i f \left(t - \frac{\hat{\Omega} \cdot \vec{x}}{v_l}\right)}, \quad (2)$$

where v_l represents the wave speed of a Love wave. Then, the two-point correlation is:

$$\langle S_H^*(f, \hat{\Omega}) S_H'(f', \hat{\Omega}') \rangle = \delta(f - f') \delta^2(\hat{\Omega}, \hat{\Omega}') H_{S,H}(f, \hat{\Omega}), \quad (3)$$

and the measurement in channel α of seismometer i would be:

$$d_{i,\alpha}(\vec{x}, t) = \vec{l}(\vec{x}, t) \cdot \hat{\alpha}, \quad (4)$$

where α can be (x, y, or z), allows us to compute the cross-correlation Y (between channel α of seismometer i and channel β of seismometer j):

$$\langle Y_{i\alpha,j\beta} \rangle = 2T\Delta f \int d\hat{\Omega} e^{-(z_i+z_j)/h} H_{S,H}(\hat{\Omega}) e_{H,\alpha}(\hat{\Omega}) e_{H,\beta}(\hat{\Omega}) e^{2\pi i f \hat{\Omega} \cdot \Delta \vec{x} / v_l}, \quad (5)$$

where the frequency dependence in Eq. 2 was ignored as in the preceding analysis and $e_{H,\alpha}(\hat{\Omega}) = \vec{e}_H(\hat{\Omega}) \cdot \hat{\alpha}$. Expanding the spatial function in some basis, e.g. pixels or spherical harmonics:

$$H_{S,H}(\hat{\Omega}) = \sum_a S_{Ha} Q_a(\hat{\Omega}), \quad (6)$$

allows us to write the γ -functions:

$$\gamma_{Ha} = \int d\hat{\Omega} Q_a(\hat{\Omega}) e_{H,\alpha}(\hat{\Omega}) e_{H,\beta}(\hat{\Omega}) e^{2\pi i f \hat{\Omega} \cdot \Delta \vec{x} / v_l}, \quad (7)$$

which, finally, enables us to write the cross-correlation as a sum over these new γ -functions:

$$\langle Y_{i\alpha,j\beta} \rangle = 2T\Delta f e^{-(z_i+z_j)/h} S_{Ha} \gamma_{Ha}. \quad (8)$$

0.1.1 Theory and Eigenfunction Approximation

Following the standard theory in seismology (Aki and Richards, 2009), SH-waves do not exist in a homogeneous halfspace, so let us consider these waves in an isotropic and vertically heterogeneous halfspace with corresponding elastic moduli that are smoothly-varying with depth. Then, assuming a plane-wave solution as above and appropriate boundary conditions (namely $l_1 \rightarrow 0$ for sufficiently large z and $\mu(z) \frac{dl_1}{dz} = 0$ at the free surface), the equation of motion takes the form:

$$-\omega^2 \rho(z) l_1 = \frac{d}{dz} \left[\mu(z) \frac{dl_1}{dz} \right] - k^2 \mu(z) l_1, \quad (9)$$

where l_1 , most-generally, is the eigenfunction that captures the displacement field's dependence on depth, frequency, and wavenumber; i.e. $l_1 = l_1(z, \omega, k)$. Equation 9 has dependence on the shear modulus, $\mu(z)$, and density, $\rho(z)$, of the medium; this must be accounted for by using an appropriate shear-wave velocity profile. By definition, the shear-wave velocity is:

$$\beta(z) = \sqrt{\frac{\mu(z)}{\rho(z)}}. \quad (10)$$

Assuming density is constant with depth, i.e. $\rho(z) = \rho_0$, we consider those profiles modelled by a power-law: $\beta(z) \sim z^{-\alpha}$. Furthermore, Haney and Tsai (2015) suggests an approximate eigenfunction solution for fundamental-mode SH-waves: $l_1 \sim e^{-akz}$. This suggests that the analysis in Sec. 0.1 can be done with a more complex eigenfunction solution, assuming the fundamental-mode of SH-waves contribute more significantly than other modes. In keeping consistent with the note, the eigenfunction in Haney and Tsai (2015) would be: $l_1 = l_1(z, \omega, k) \sim l_1(z, f, v_l) = e^{2\pi a \frac{fz}{v_l}}$ if $k = \omega/v_l$ and $\omega = 2\pi/f$. Note its dependence on Love wave velocity; this value is usually obtained experimentally. Scanning through different values of attenuation factors, $\alpha \in [0.250 \ 0.275 \ 0.300 \ 0.325 \ 0.350 \ 0.375 \ 0.400]$, they found $a = 0.85 \pm 0.09$.

References

- Keiti Aki and Paul G. Richards. *Quantitative Seismology*. University Science Books, Mill Valley, CA, 2nd edition, 2009.
- Matthew M. Haney and Victor C. Tsai. Nonperturbational surface-wave inversion: A Dix-type relation for surface waves. *Geophysics*, 80(6):EN167–EN177, 2015. URL doi:10.1190/geo2014-0612.1.
- Thorne Lay and Terry C. Wallace. *Modern Global Seismology*, volume 58 of *International Geophysics Series*. Academic Press, San Diego, CA, 1995.